## Characteristics and Classification of Living Organisms

## **Question Paper 4**

Level	IGCSE	
Subject	Biology	
Exam Board	CIE	
Topic	Characteristics and Classification of Living Organisms	
Paper Type	(Extended) Theory Paper	
Booklet	Question Paper 4	

Time Allowed: 68 minutes

Score: /56

Percentage: /100

## 1 Fig. 1.1 shows seven marsupial mammals.



not drawn to scale

Fig. 1.1

(a) tail visible go to 2 (b) no tail visible go to 3 (a) back feet at least twice as long as front feet go to 4 (b) back feet and front feet of similar length go to 5 (a) large ears relative to the size of the head Phascolarctos cinereus (b) small ears relative to the size of the head Vombatus ursinus (a) tail at least twice as long as body Sminthopsis longicaudata (b) tail less than twice as long as body Macropus rufus (a) uniform body colouring Paljara tirarense (b) markings on body go to 6 (a) white band across back and chest Sarcophilus harrisii (b) no white band across back and chest Dasyurus maculatus G  b) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a gamete and a female gamete.  (i) Name the process that results in the formation of haploid gametes.	(a) tail visible go to 2  (b) no tail visible go to 3  (a) back feet at least twice as long as front feet go to 4  (b) back feet and front feet of similar length go to 5  (a) large ears relative to the size of the head Phascolarctos cinereus  (b) small ears relative to the size of the head Vombatus ursinus  (a) tail at least twice as long as body Sminthopsis longicaudata  (b) tail less than twice as long as body Macropus rufus  (a) uniform body colouring Paljara tirarense  (b) markings on body go to 6  (a) white band across back and chest Sarcophilus harrisii  (b) no white band across back and chest Dasyurus maculatus G  (b) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a gamete and a female gamete.	(ii)	(ii) Use the key to identify each species. Write the letter of each species (A to G) in the correct box beside the key. One has been done for you.					
(b) no tail visible go to 3  (a) back feet at least twice as long as front feet go to 4  (b) back feet and front feet of similar length go to 5  (a) large ears relative to the size of the head Phascolarctos cinereus  (b) small ears relative to the size of the head Vombatus ursinus  (a) tail at least twice as long as body Sminthopsis longicaudata  (b) tail less than twice as long as body Macropus rufus  (a) uniform body colouring Paljara tirarense  (b) markings on body go to 6  (a) white band across back and chest Sarcophilus harrisii  (b) no white band across back and chest Dasyurus maculatus G  (b) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a gamete and a female gamete.	(b) no tail visible go to 3  (a) back feet at least twice as long as front feet go to 4  (b) back feet and front feet of similar length go to 5  (a) large ears relative to the size of the head Phascolarctos cinereus  (b) small ears relative to the size of the head Vombatus ursinus  (a) tail at least twice as long as body Sminthopsis longicaudata  (b) tail less than twice as long as body Macropus rufus  (a) uniform body colouring Paljara tirarense  (b) markings on body go to 6  (a) white band across back and chest Sarcophilus harrisii  (b) no white band across back and chest Dasyurus maculatus G  (b) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a regamete and a female gamete.  (i) Name the process that results in the formation of haploid gametes.	(2)	-	go to 2				
(a) back feet at least twice as long as front feet (b) back feet and front feet of similar length (a) large ears relative to the size of the head (b) small ears relative to the size of the head (c) small ears relative to the size of the head (d) tail at least twice as long as body (e) tail less than twice as long as body (f) markings on body (g) markings on body (g) white band across back and chest (h) no white band across back and chest (g) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a gamete and a female gamete.	(a) back feet at least twice as long as front feet (b) back feet and front feet of similar length (a) large ears relative to the size of the head (b) small ears relative to the size of the head (c) small ears relative to the size of the head (d) tail at least twice as long as body (e) tail less than twice as long as body (f) markings on body (g) to 6 (h) markings on body (g) to 6 (h) white band across back and chest (h) no white band across back and chest (h) Name the process that results in the formation of haploid gametes.							
(b) back feet and front feet of similar length go to 5  (a) large ears relative to the size of the head Phascolarctos cinereus  (b) small ears relative to the size of the head Vombatus ursinus  (a) tail at least twice as long as body Sminthopsis longicaudata  (b) tail less than twice as long as body Macropus rufus  (a) uniform body colouring Paljara tirarense  (b) markings on body go to 6  (a) white band across back and chest Sarcophilus harrisii  (b) no white band across back and chest Dasyurus maculatus G  (b) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a gamete and a female gamete.	(b) back feet and front feet of similar length go to 5  (a) large ears relative to the size of the head Phascolarctos cinereus  (b) small ears relative to the size of the head Vombatus ursinus  (a) tail at least twice as long as body Sminthopsis longicaudata  (b) tail less than twice as long as body Macropus rufus  (a) uniform body colouring Paljara tirarense  (b) markings on body go to 6  (a) white band across back and chest Sarcophilus harrisii  (b) no white band across back and chest Dasyurus maculatus G  De) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a regamete and a female gamete.  (i) Name the process that results in the formation of haploid gametes.							
(a) large ears relative to the size of the head  (b) small ears relative to the size of the head  (a) tail at least twice as long as body  (b) tail less than twice as long as body  (a) uniform body colouring  (b) markings on body  (c) markings on body  (d) white band across back and chest  (e) no white band across back and chest  (f) Dasyurus maculatus  (g) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a gamete and a female gamete.	(a) large ears relative to the size of the head Phascolarctos cinereus  (b) small ears relative to the size of the head Vombatus ursinus  (a) tail at least twice as long as body Sminthopsis longicaudata  (b) tail less than twice as long as body Macropus rufus  (a) uniform body colouring Paljara tirarense  (b) markings on body go to 6  (a) white band across back and chest Sarcophilus harrisii  (b) no white band across back and chest Dasyurus maculatus G  (b) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a regamete and a female gamete.  (i) Name the process that results in the formation of haploid gametes.							
(b) small ears relative to the size of the head   (a) tail at least twice as long as body   (b) tail less than twice as long as body   (a) uniform body colouring   (b) markings on body   (c) markings on body   (d) white band across back and chest   (e) no white band across back and chest   (f) Dasyurus maculatus   (g) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a gamete and a female gamete.	(b) small ears relative to the size of the head Vombatus ursinus  (a) tail at least twice as long as body Sminthopsis longicaudata  (b) tail less than twice as long as body Macropus rufus  (a) uniform body colouring Paljara tirarense  (b) markings on body go to 6  (a) white band across back and chest Sarcophilus harrisii  (b) no white band across back and chest Dasyurus maculatus G  (b) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a regamete and a female gamete.  (i) Name the process that results in the formation of haploid gametes.							
(a) tail at least twice as long as body  (b) tail less than twice as long as body  (a) uniform body colouring  (b) markings on body  (c) white band across back and chest  (d) no white band across back and chest  (e) no white band across back and chest  (f) Dasyurus maculatus  (g) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a gamete and a female gamete.	(a) tail at least twice as long as body  (b) tail less than twice as long as body  (a) uniform body colouring  (b) markings on body  (c) white band across back and chest  (d) no white band across back and chest  (e) no white band across back and chest  (f) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a regamete and a female gamete.  (i) Name the process that results in the formation of haploid gametes.							
(b) tail less than twice as long as body  (a) uniform body colouring  (b) markings on body  (c) white band across back and chest  (d) no white band across back and chest  (e) no white band across back and chest  (f) Dasyurus maculatus  (g) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a gamete and a female gamete.	(b) tail less than twice as long as body  (a) uniform body colouring  (b) markings on body  (c) white band across back and chest  (d) no white band across back and chest  (e) no white band across back and chest  (f) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a gamete and a female gamete.  (i) Name the process that results in the formation of haploid gametes.		tail at least twice as long as body	Sminthopsis longicaudata				
(b) markings on body  (a) white band across back and chest  (b) no white band across back and chest  (c) Dasyurus maculatus  (d) Dasyurus maculatus  (e) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a gamete and a female gamete.	(b) markings on body (a) white band across back and chest (b) no white band across back and chest (b) no white band across back and chest Dasyurus maculatus  G  Dispute the production occurs in all mammals. A zygote is formed from the fertilisation of a regamete and a female gamete.  (i) Name the process that results in the formation of haploid gametes.							
(a) white band across back and chest  (b) no white band across back and chest  Dasyurus maculatus  G  Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a gamete and a female gamete.	(a) white band across back and chest  (b) no white band across back and chest  Dasyurus maculatus  G  Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a gamete and a female gamete.  (i) Name the process that results in the formation of haploid gametes.	(a)	uniform body colouring	Paljara tirarense				
(b) no white band across back and chest Dasyurus maculatus G  b) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a gamete and a female gamete.	(b) no white band across back and chest  Dasyurus maculatus  G  Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a gamete and a female gamete.  (i) Name the process that results in the formation of haploid gametes.	(b)	markings on body	go to 6				
b) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a gamete and a female gamete.	<ul> <li>b) Sexual reproduction occurs in all mammals. A zygote is formed from the fertilisation of a gamete and a female gamete.</li> <li>(i) Name the process that results in the formation of haploid gametes.</li> </ul>	(a)	white band across back and chest	Sarcophilus harrisii				
gamete and a female gamete.	gamete and a female gamete.  (i) Name the process that results in the formation of haploid gametes.	(b)	no white band across back and chest	and chest Dasyurus maculatus G				
		ga	amete and a female gamete.	on of haploid gametes.				

.....[3]

(c)	Marsupials differ from other mammals by giving birth to relatively undeveloped of Female humans have a placenta and therefore give birth to more developed offspring			
	(i)	Describe the role of the placenta in humans.		
		[4]		
	(ii)	In humans, the placenta is connected to the amniotic sac which contains amniotic fluid.		
		State <b>two</b> functions of the amniotic fluid.		
		[2]		
		[Total: 14]		

Molluscs are important animals in many aquatic and terrestrial ecosystems. 2

Fig. 1.1 shows four species of mollusc that live in the sea.




Fig. 1.1

(a) State two features shown by all mollusc species.

1	
2	

(b) State two features, visible in Fig. 1.1, in which the octopus differs from the other three molluscs.

1	
2	[2]

.....

c)	The edible mussel, <i>Mytilus edulis</i> , is attached to rocks that are exposed to the air at low tide.
	Use Fig. 1.1 to suggest how an edible mussel is adapted to attach to rocks and survive when exposed to the air.
	[2]
d)	The zebra mussel, <i>Dreissena polymorpha</i> , is a freshwater mussel that originates from rivers in southern Russia.
	The mussel was introduced into the Great Lakes of North America and has increased in huge numbers with serious effects on the food webs of the lakes.
	Explain why an introduced species, such as the zebra mussel, can have serious effects on the populations of the species that are already living in the area.
	—— <b>ТТПТТПОМ</b> —— [3]

(e)	The freshwater mussel, <i>Pletholophus swinhoei</i> , was used in a project to monitor water pollution by chemical waste in northern Vietnam.
	This was done by regularly counting the number of mussels in the river.
	Suggest the advantages of using freshwater mussels to monitor the pollution of water instead of carrying out chemical analysis of the water.
	[2]
(f)	Non-biodegradable plastics are a serious problem in many aquatic ecosystems.
	Explain the harm that non-biodegradable plastics may cause to organisms in aquatic ecosystems.
	[3]
	[Total: 14]

Fig. 1.1 shows a flowering shoot of tiger lily, *Lilium tigrinum*.




Fig. 1.1

		[1]
 	 ······	

(b) Name the parts labelled A to D.

(a) State the name of the genus of the tiger lily.

A	
В	
С	
D	[4]

(c)	The tiger lily plant is a monoc	otyledon.	
	List two features, visible in F	ig. 1.1, that show it is a mono	ocotyledon.
	1		
	2		[2]
(d)	The tiger lily in Fig. 1.1 reprod		
	Plants reproduce sexually an	d asexually.	
	Complete Table 1.1 to show reproduction to a flowering pl		ntages of asexual and sexual
		Table 1.1	
	type of reproduction in flowering plants	advantages	dis
	asexual		
	CHEMI	STRYON	LINE
	sexual	IIITION	
		OTITOIN	
			[4]

[Total: 11]

4 Reed warblers are small birds that migrate over long distances between western Africa and northern Europe.

Fig. 5.1 shows a reed warbler, Acrocephalus scirpaceus.



Fig. 5.1

(a) State three characteristic features of birds that are visible in Fig. 5.1.

1	
2	
3	[3]
•	[~]

A study was carried out in Sweden into the effects of natural selection on wing length in reed warblers.

The wings of young reed warblers reach their maximum length a few days after leaving the nest.

At this age the wing length in millimetres of each bird was recorded. Each bird was identified by putting a small ring around one of its legs.

When the birds were caught in net traps as adults, the information on the rings was used to identify specific birds and their ages.

The length of time between ringing and trapping was recorded for each bird that was identified before it was released.

The mean age at trapping was calculated for birds with each wing length.

The results are shown in Table 5.1.

Table 5.1

wing length at ringing / mm	number of birds trapped	mean age at trapping / days
63 or less	24	253
64	72	256
65	1	297
66	1	346
67	1	349
68	1	270
69	66	237
70 or more	23	199
	total = 771	

(b) (i)	Explain why wing length is an example of continuous variation.
	[2
(ii)	Suggest a feature of reed warblers, <b>other than wing length</b> , that shows continuous variation.
	[1]

The researchers concluded that reed warblers with a wing length of 66-67 mm had the best chance of survival.		
(i)	Describe the evidence from Table 5.1 that supports this conclusion.	
	[4]	
(ii)	The researchers also suggested that more evidence was needed to make this conclusion.	
	Suggest what other evidence would show that birds with wings 66-67 mm in length have the best chance of survival.	
	CHEMISTRY ONLINE	
	—— TIIITION —— [3]	
	[၁]	

(c)

(d)	Scientists have discovered that genes are responsible for wing length in reed warblers. The most common length of wing has been 66-67 mm for many generations of these
	birds.  Explain how natural selection may be responsible for maintaining the mean wing length of reed warblers at 66-67 mm.
	[4]
	[Total: 17]