

Inheritance

Mark Scheme 5

Level	IGCSE
Subject	Biology
Exam Board	CIE
Topic	Inheritance
Paper Type	(Extended) Theory Paper
Booklet	Mark Scheme 5

CHEMISTRY ONLINE
— TUITION —

Time Allowed: 61 minutes

Score: /51

Percentage: /100

Question	Answers	Marks	Guidance
1 (a)	<p><i>general marks</i> roots absorb water ; idea of <u>both</u> gaining water over a large, volume / area, of soil ; AVP ;</p> <p>A has deep roots / go a long way down ; to gain water that drains through soil / reach water table / AW ;</p> <p>B has shallow roots / wide spreading roots / AW ; absorbs water, before it drains <i>or</i> evaporates / immediately after rainfall ;</p>	[max 4]	<p>NB water absorption and area marks given once only</p> <p>R long roots unqualified</p>
(b)	<p>thick cuticle ; longer distance for diffusion / not easy for water to pass through / ref to impermeable ;</p> <p>rolled leaves ; air trapped inside rolled leaf has <u>higher</u> humidity AW / stomata protected from wind <i>or</i> moving air (reduces transpiration) ;</p> <p>sunken stomata / stomata in pits <i>or</i> grooves <i>or</i> depressions ; chamber has <u>higher</u> humidity AW / stomata protected from wind <i>or</i> moving air (so reducing transpiration) ;</p> <p>hairs on leaf ; reduce air flow over the surface (so reducing transpiration) / increase humidity by 'trapping' water (molecules) ;</p> <p>small leaves / leaves reduced to spines / leaves are needles / no leaves / leaves shed in very dry periods ; small(er) / no surface area (for transpiration) ;</p> <p>fewer stomata / stomata closed during hot parts of day ; stomata are pores through which water can pass (so reducing transpiration) ;</p>	[2 + 2]	<p>R cuticle unqualified or ref to 'waxy' without description of thickness</p> <p>Must be TWO descriptions (max) with appropriate linked explanations</p> <p>explanations alone cannot be accepted</p> <p>A correct references to water potential / concentration gradient for rolled leaves or sunken stomata</p> <p>IGNORE references to succulent leaves and storage (not water loss)</p> <p>'sharp' leaves also need to be small</p>

Question	E	Answers	Marks	Guidance																		
1 (c)		<table><tr><th>tissue</th><th>substances transported</th><th>source</th><th>sink</th></tr><tr><td>xylem</td><td>water, ions / named ion / mineral / salts ;</td><td>roots ;</td><td>stem / growing points / buds / leaf / flower / fruit / seed / storage organ ;</td></tr><tr><td rowspan="4">phloem</td><td rowspan="4">Sucrose / sugar, amino acids ;</td><td colspan="2"><i>either</i></td></tr><tr><td>leaf ;</td><td>stem / growing points / buds / root / flower / fruit / seed / storage organ ;</td></tr><tr><td colspan="2"><i>or</i></td></tr><tr><td>storage organ ;</td><td><u>young AW</u> leaf / stem / growing points / buds / root ;</td></tr></table>	tissue	substances transported	source	sink	xylem	water, ions / named ion / mineral / salts ;	roots ;	stem / growing points / buds / leaf / flower / fruit / seed / storage organ ;	phloem	Sucrose / sugar, amino acids ;	<i>either</i>		leaf ;	stem / growing points / buds / root / flower / fruit / seed / storage organ ;	<i>or</i>		storage organ ;	<u>young AW</u> leaf / stem / growing points / buds / root ;	[6]	<p>NB substances transported score:-</p> <p>ONE mark for TWO correct responses</p> <p>R references to single cells as sources or sinks e.g. root hairs</p> <p>R glucose</p> <p>mark each box independently</p>
tissue	substances transported	source	sink																			
xylem	water, ions / named ion / mineral / salts ;	roots ;	stem / growing points / buds / leaf / flower / fruit / seed / storage organ ;																			
phloem	Sucrose / sugar, amino acids ;	<i>either</i>																				
		leaf ;	stem / growing points / buds / root / flower / fruit / seed / storage organ ;																			
		<i>or</i>																				
		storage organ ;	<u>young AW</u> leaf / stem / growing points / buds / root ;																			
			[Total: 14]																			

Question	Answers	Marks	Additional Guidance
2 (a)	pollen transferred from, anther / stamen, to stigma ; within same <u>flower</u> / between <u>flowers</u> on same plant ; R if only 'same plant'	[2]	R complete answers given in context of fertilisation R 'single parent'
(b)	<p><i>cross 1</i></p> $\begin{array}{c} I^R I^R \times I^W I^W \\ I^R + I^W \\ I^R I^W \end{array} ;$ <p><i>cross 2</i></p> $I^R I^W \times I^R I^W$ $I^R, I^W + I^R, I^W ;$ $I^R I^R, I^R I^W, (I^R I^W), I^W I^W ;$ <p>1 <u>red</u> : 2 <u>pink</u> : 1 <u>white</u> ; A 25% red : 50% pink : 25% white A multiples, e.g. 2 red: 4 pink : 2 white</p> <p>R if two different ratios given</p>	[4]	<p>A other notation, e.g. R and r or mixture, e.g. I^R and W. R I^{RR}, etc.</p> <p><i>cross 1</i> 1 mark for parental genotypes, gametes and offspring all correct. Any mistake and no mark awarded.</p> <p><i>cross 2</i> 1 mark for cross genotypes and gametes all correct. Any mistake and no mark awarded.</p> <p>1 mark for giving all three genotypes (on answer line or in the white space e.g. in Punnett square). If correct on answer line ignore any errors in working.</p> <p>1 mark for ratio of offspring phenotypes and colours R if no colours given</p>
(c)	$I^R I^W \times I^W I^W$ $I^R, I^W + I^W ;$ $I^R I^W, I^W I^W ;$ <p>1 (pink) : 1 (white) ; R if two different ratios given</p>	[3]	<p>1 mark for parental genotypes and gametes all correct. Any mistake and no mark awarded.</p> <p>1 mark for offspring genotypes</p> <p>1 mark for ratio (colours not necessary) A if no colours given</p>

Question	E	Answers	Marks	Additional Guidance
2 (d)	1 2 3 4 5 6 7 8 9 10	ref. to meiosis ; mutation can occur <u>in meiosis</u> ; (gives) variation / diversity ; R 'varied species (plural)' ref. to, alleles / genes / DNA, from different, plants / parents ; allows mutations to be, expressed / AW ; allows adaptation to, new conditions / changed environment / AW ; (new species) can evolve / allows natural selection to occur ; seeds are dispersed ; R dispersed unqualified, R pollen dispersal can colonise new areas / AW ; less competition (with parent plant / among offspring) ;	 [max 4]	R sexual reproduction allows mutations to occur A may allow resistance to disease A 'suited to' / survive / AW for adapted R 'passed on by natural selection' R 'new species are made' A 'go to new areas' or 'spread to new areas' <i>competition is in context of seed dispersal not pollen dispersal</i> R 'multiply quicker'
			[Total: 13]	

3 (a) (length of) DNA / part of chromosome / on a chromosome ,
that codes for a protein or polypeptide or enzyme / controls a characteristic ; [1]

(b) $H^N H^S \times H^N H^S$; accept N and S

$H^N, H^S + H^N, H^S$; gametes must be clear *accept on dotted line or in Punnett square*

$H^S H^S$; *ecf from correct gametes if wrong parental genotype* [3]

(c) check <http://www.sicklecellsociety.org/education/healthpr.htm> for AVPs

- 1 red (blood) cells become, sickle shaped / distorted / AW ; **R** abnormal unqualified
- 2 in areas of low oxygen concentrations / in tissues ;
- 3 fewer / less elastic / less flexible / short-lived, red blood cells ; *ora*
- 4 less haemoglobin ;
- 5 blood / haemoglobin, less efficient at transporting oxygen ; **R** no oxygen
- 6 less respiration ; **R** no respiration
- 7 less energy / fatigued / exhaustion / less active / feeling faint or tired / breathless ;
- 8 capillaries are blocked ;
- 9 pain ;
- 10 death of tissues linked to blood supply ;
- 11 'sickle cell crisis' ; **A** 'attacks needing oxygen'
- 12 slow / poor, growth ;
- 13 susceptible to infections ;
- 14 reduced life span ;
- 15 AVP ;
- 16 AVP ;

[4 max]

- 3 (d) 1 *idea that* areas with high percentage of sickle cell (allele) are places with malaria ;
 2 $H^S H^S$ / homozygous recessive, reduced life span because of sickle cell anaemia ;
 3 $H^N H^N$ / homozygous dominant / without H^S , susceptible to malaria / AW ;
 4 $H^N H^S$ / heterozygous / carrier/ with H^S , resistant / not affected / less susceptible ;
 A $H^S H^S$ **R** immune / immunity
 5 $H^N H^S$ (carrier) survive and have children / $H^N H^N$ or $H^S H^S$ do not ;
 6 $H^N H^S$ / carrier, pass on the allele / H^S ;
 7 (if $H^N H^S \times H^N H^S$) 1 in 4 chance of, $H^S H^S$ / homozygous recessive ;
 8 2 in 4 / 50% / $\frac{1}{2}$, have advantage of resistance to malaria ; [5 max]

- (e) 1 *idea that* distinct groups / categories ; ref to bar chart
 2 *either* sickle cell anaemia ($H^S H^S$), sickle cell trait ($H^N H^S$), normal ($H^N H^N$) / or normal, anaemic ; **A** 'some people have disease, some do not'
 A 'some people have the allele, some do not'
 3 no intermediates / no continuous scale of anaemia / AW ;
 4 genetic condition / environment has no effect (or its expression) ;
 A ref to small number of, genes / alleles, involved [3 max]

[Total: 16]

CHEMISTRY ONLINE
 — TUITION —

4 (i)

process	materials moved	source of materials in the plant	sink for materials in the plant
transpiration	water + (mineral) salts / AW ; A ions / minerals / named ion R nutrients	roots / root hairs ;	leaves / shoot / stem ; A flowers / fruits named, cell(s) / tissue(s)
translocation	<i>two from</i> sugars / sucrose amino acids ions / minerals / AW hormones / named hormone; R glucose R nutrients	leaves / (named) storage organ / seed(s) / cotyledon ;	roots / stem / shoot / named growing region / (named) storage organ ; A buds / flowers / fruits / tubers A named cell(s) / tissue(s)

[6]

(ii) *answer needs to make clear which structures are source and sink*

during germination / AW, (source is) seed / cotyledon ;
idea that leaves grow and start to photosynthesise (so become source) ;

leaves may, be shed / die / be shaded / AW ;
leaves may stop photosynthesising (so become sink) / AW ; **A** 'slow down'

(in early growth) root (is sink) ;
(later) flowers / fruits / seeds / tubers / AW (become sinks) ;

[max. 2]

[Total: 8]