

# Electrolysis, Electrode Potentials & Cells

## Question Paper 2

<b>Level</b>	International A Level
<b>Subject</b>	Chemistry
<b>Exam Board</b>	CIE
<b>Topic</b>	Electrochemistry
<b>Sub-Topic</b>	Electrolysis, Electrode Potentials & Cells
<b>Paper Type</b>	Theory
<b>Booklet</b>	Question Paper 2

**Time Allowed:** 80 minutes

**Score:** /66

**Percentage:** /100

**Grade Boundaries:**

A*	A	B	C	D	E	U
>85%	77.5%	70%	62.5%	57.5%	45%	<45%

1 (a) (i) With the aid of a fully-labelled diagram, describe the standard hydrogen electrode.

(ii) Use the *Data Booklet* to calculate the standard cell potential for the reaction between  $\text{Cr}^{2+}$  ions and  $\text{Cr}_2\text{O}_7^{2-}$  ions in acid solution, and construct a balanced equation for the reaction.

$$E_{\text{cell}}^{\ominus} = \dots\dots\dots \text{V}$$

equation .....

(iii) Describe what you would see if a blue solution of  $\text{Cr}^{2+}$  ions was added to an acidified solution of  $\text{Cr}_2\text{O}_7^{2-}$  ions until reaction was complete.

.....  
.....

[8]

CHEMISTRY ONLINE  
— TUITION —

- (b) A buffer solution is to be made using  $1.00 \text{ mol dm}^{-3}$  ethanoic acid,  $\text{CH}_3\text{CO}_2\text{H}$ , and  $1.00 \text{ mol dm}^{-3}$  sodium ethanoate,  $\text{CH}_3\text{CO}_2\text{Na}$ .

Calculate to the nearest  $1 \text{ cm}^3$  the volumes of each solution that would be required to make  $100 \text{ cm}^3$  of a buffer solution with pH 5.50.

Clearly show all steps in your working.

$$K_a(\text{CH}_3\text{CO}_2\text{H}) = 1.79 \times 10^{-5} \text{ mol dm}^{-3}$$

volume of  $1.00 \text{ mol dm}^{-3} \text{ CH}_3\text{CO}_2\text{H} = \dots\dots\dots \text{ cm}^3$


volume of  $1.00 \text{ mol dm}^{-3} \text{ CH}_3\text{CO}_2\text{Na} = \dots\dots\dots \text{ cm}^3$   
[4]

- (c) Write an equation to show the reaction of this buffer solution with each of the following.

(i) added  $\text{HCl}$  .....

(ii) added  $\text{NaOH}$  .....  
[2]

- (d) Choose **one** reaction in organic chemistry that is catalysed by an acid, and write the structural formulae of the reactants and products in the boxes below.



[3]


[Total: 17]

2 (a) What is meant by the term *standard electrode potential*, SEP?

.....  
.....


[2]

(b) Draw a fully labelled diagram of the apparatus you could use to measure the SEP of the Fe<sup>3+</sup>/Fe<sup>2+</sup> electrode.



[5]

(c) The reaction between Fe<sup>3+</sup> ions and I<sup>-</sup> ions is an equilibrium reaction.



(i) Use the *Data Booklet* to calculate the  $E_{\text{cell}}^{\ominus}$

.....

(ii) Hence state, with a reason, whether there will be more products or more reactants at equilibrium.

.....  
.....

(iii) Write the expression for  $K_c$  for this reaction, and state its units.

$K_c =$

units .....

An experiment was carried out using solutions of  $\text{Fe}^{3+}(\text{aq})$  and  $\text{I}^{-}(\text{aq})$  of equal concentrations.  $100\text{ cm}^3$  of each solution were mixed together, and allowed to reach equilibrium.

The concentrations at equilibrium of  $\text{Fe}^{3+}(\text{aq})$  and  $\text{I}_2(\text{aq})$  were as follows.

$$[\text{Fe}^{3+}(\text{aq})] = 2.0 \times 10^{-4} \text{ mol dm}^{-3}$$

$$[\text{I}_2(\text{aq})] = 1.0 \times 10^{-2} \text{ mol dm}^{-3}$$

- (iv) Use these data, together with the equation given in (c), to calculate the concentrations of  $\text{Fe}^{2+}(\text{aq})$  and  $\text{I}^{-}(\text{aq})$  at equilibrium.

$$[\text{Fe}^{2+}(\text{aq})] = \dots\dots\dots \text{ mol dm}^{-3}$$

$$[\text{I}^{-}(\text{aq})] = \dots\dots\dots \text{ mol dm}^{-3}$$

- (v) Calculate the  $K_c$  for this reaction.

$$K_c = \dots\dots\dots$$

[8]

[Total: 15]

3 Chlorine gas is manufactured by the electrolysis of brine using a diaphragm cell.

(a) (i) Write half-equations, including state symbols, for the reactions occurring at **each** of the electrodes of a diaphragm cell.

anode .....

cathode .....

(ii) In the diaphragm cell, the anode is made of titanium and the cathode is made of steel.

Suggest why steel is never used for the anode.

.....  
.....

[3]

(b) Chlorine is very reactive and will form compounds by direct combination with many elements.

Describe what you would see when chlorine is passed over separate heated samples of sodium and phosphorus.

In **each** case write an equation for the reaction.

sodium

.....  
.....  
.....

phosphorus

.....  
.....  
.....

[4]

- (c) Chlorine reacts with aqueous sodium hydroxide in two different ways, depending on the conditions used. In each case, water, sodium chloride and one other chlorine-containing compound are formed.

For **each** condition below, give the formula of the **other** chlorine-containing compound and state the oxidation number of chlorine in it.

condition	formula of <b>other</b> chlorine-containing compound	oxidation number of chlorine in this compound
cold dilute NaOH(aq)		
hot concentrated NaOH(aq)		

[4]

- (d) Magnesium chloride,  $MgCl_2$ , and silicon tetrachloride,  $SiCl_4$ , each dissolve in or react with water.

Suggest the approximate pH of the solution formed in **each** case.

$MgCl_2$  .....  $SiCl_4$  .....

Explain, with the aid of an equation, the difference between the two values.

.....

.....

.....


.....

..... [5]

CHEMISTRY ONLINE  
— TUITION —

[Total: 16]

- 4 (a) The diagram below shows an incomplete experimental set-up needed to measure the  $E_{\text{cell}}$  of a cell composed of the standard  $\text{Cu}^{2+}/\text{Cu}$  electrode and an  $\text{Ag}^+/\text{Ag}$  electrode.



- (i) State the chemical composition of  
 solution A, .....
- electrode B. ....
- (ii) Complete the diagram to show the whole experimental set-up. [4]

- (b) The above cell is not under standard conditions, because the  $[\text{Ag}^+]$  in a saturated solution of  $\text{AgCl}$  is much less than  $1.0 \text{ mol dm}^{-3}$ . The  $E_{\text{electrode}}$  is related to  $[\text{Ag}^+]$  by the following equation.

equation 1 
$$E_{\text{electrode}} = E_{\text{electrode}}^{\ominus} + 0.06 \log[\text{Ag}^+]$$

- (i) Use the *Data Booklet* to calculate the  $E_{\text{cell}}^{\ominus}$  if the cell was operating under standard conditions.

CHEMISTRY ONLINE  
— TUITION —

$E_{\text{cell}}^{\ominus} = \dots\dots\dots \text{V}$

In the above experiment, the  $E_{\text{cell}}$  was measured at +0.17V.

- (ii) Calculate the value of  $E_{\text{electrode}}$  for the  $\text{Ag}^+/\text{Ag}$  electrode in this experiment.  
 .....
- (iii) Use equation 1 to calculate  $[\text{Ag}^+]$  in the saturated solution.

$[\text{Ag}^+] = \dots\dots\dots \text{mol dm}^{-3}$

[3]



