

# Uniform Electric Fields

## Question paper 2

<b>Level</b>	International A Level
<b>Subject</b>	Physics
<b>Exam Board</b>	CIE
<b>Topic</b>	Electric Fields
<b>Sub Topic</b>	Uniform Electric Fields
<b>Paper Type</b>	Theory
<b>Booklet</b>	Question paper 2

**Time Allowed:** 56 minutes

**Score:** /46

**Percentage:** /100


CHEMISTRY ONLINE

A*	A	B	C	D	E	U
>85%	77.5%	70%	62.5%	57.5%	45%	<45%

- 1 (a) Define *electric field strength*.

.....  
.....[1]

- (b) Two horizontal metal plates are 20 mm apart in a vacuum. A potential difference of 1.5 kV is applied across the plates, as shown in Fig. 4.1.



**Fig. 4.1**

A charged oil drop of mass  $5.0 \times 10^{-15}$  kg is held stationary by the electric field.

- (i) On Fig. 4.1, draw lines to represent the electric field between the plates. [2]  
(ii) Calculate the electric field strength between the plates.

electric field strength = .....  $\text{V m}^{-1}$  [1]

- (iii) Calculate the charge on the drop.

charge = ..... C [4]

- (iv) The potential of the upper plate is increased. Describe and explain the subsequent motion of the drop.

.....  
.....  
.....[2]

- 2 Two horizontal metal plates are separated by distance  $d$  in a vacuum. A potential difference  $V$  is applied across the plates, as shown in Fig. 6.1.




Fig. 6.1

A horizontal beam of  $\alpha$ -particles from a radioactive source is made to pass between the plates.

- (a) State and explain the effect on the deflection of the  $\alpha$ -particles for each of the following changes:

- (i) The magnitude of  $V$  is increased.

.....  
..... [1]

- (ii) The separation  $d$  of the plates is decreased.

.....  
..... [1]

- (b) The source of  $\alpha$ -particles is replaced with a source of  $\beta$ -particles.  
Compare, with a reason in each case, the effect of each of the following properties on the deflections of  $\alpha$ - and  $\beta$ -particles in a uniform electric field:

(i) charge

.....  
.....  
..... [2]

(ii) mass

.....  
.....  
..... [2]

(iii) speed


.....  
.....  
..... [1]

- (c) The electric field gives rise to an acceleration of the  $\alpha$ -particles and the  $\beta$ -particles.  
Determine the ratio

$$\frac{\text{acceleration of the } \alpha\text{-particles}}{\text{acceleration of the } \beta\text{-particles}} .$$

ratio = ..... [3]

- 3 Two oppositely-charged parallel metal plates are situated in a vacuum, as shown in Fig.



**Fig. 7.1**

The plates have length  $L$ .

The uniform electric field between the plates has magnitude  $E$ . The electric field outside the plates is zero.

A positively-charged particle has mass  $m$  and charge  $+q$ . Before the particle reaches the region between the plates, it is travelling with speed  $v$  parallel to the plates. The particle passes between the plates and into the region beyond them.

- (a) (i) On Fig. 7.1, draw the path of the particle between the plates and beyond them. [2]

- (ii) For the particle in the region between the plates, state expressions, in terms of  $E$ ,  $m$ ,  $q$ ,  $v$  and  $L$ , as appropriate, for

1. the force  $F$  on the particle,

..... [1]

2. the time  $t$  for the particle to cross the region between the plates.

..... [1]

**(b) (i)** State the law of conservation of linear momentum.

.....  
.....  
..... [2]

**(ii)** Use your answers in **(a)(ii)** to state an expression for the change in momentum of the particle.

..... [1]

**(iii)** Suggest and explain whether the law of conservation of linear momentum applies to the particle moving between the plates.


.....  
.....  
..... [2]

CHEMISTRY ONLINE  
— TUITION —

- 4 (a) State what is meant by an *electric field*.

.....  
.....[1]


- (b) The electric field between an earthed metal plate and two charged metal spheres is illustrated in Fig. 5.1.



**Fig. 5.1**

- (i) On Fig. 5.1, label each sphere with (+) or (–) to show its charge. [1]
- (ii) On Fig. 5.1, mark a region where the magnitude of the electric field is
1. constant (label this region C), [1]
  2. decreasing (label this region D). [1]

- (c) A molecule has its centre P of positive charge situated a distance of  $2.8 \times 10^{-10} \text{ m}$  from its centre N of negative charge, as illustrated in Fig. 5.2.



**Fig. 5.2**


The molecule is situated in a uniform electric field of field strength  $5.0 \times 10^4 \text{ V m}^{-1}$ . The axis NP of the molecule is at an angle of  $30^\circ$  to this uniform applied electric field. The magnitude of the charge at P and at N is  $1.6 \times 10^{-19} \text{ C}$ .

- (i) On Fig. 5.2, draw an arrow at P and an arrow at N to show the directions of the forces due to the applied electric field at each of these points. [1]
- (ii) Calculate the torque on the molecule produced by the forces in (i).

torque = ..... N m [2]



- 5 Two vertical parallel metal plates are situated 2.50 cm apart in a vacuum. The potential difference between the plates is 350 V, as shown in Fig. 6.1.



**Fig. 6.1**

An electron is initially at rest close to the negative plate and in the uniform electric field between the plates.

- (a) (i) Calculate the magnitude of the electric field between the plates.

electric field strength = .....  $\text{NC}^{-1}$  [2]

- (ii) Show that the force on the electron due to the electric field is  $2.24 \times 10^{-15} \text{ N}$ .

[2]

(b) The electron accelerates horizontally across the space between the plates. Determine

(i) the horizontal acceleration of the electron,

acceleration = .....  $\text{ms}^{-2}$  [2]

(ii) the time to travel the horizontal distance of 2.50 cm between the plates.

time = ..... s [2]

(c) Explain why gravitational effects on the electron need not be taken into consideration in your calculation in (b).

.....  
.....  
..... [2]