

# Nuclear Physics

## Question paper 3

<b>Level</b>	International A Level
<b>Subject</b>	Physics
<b>Exam Board</b>	CIE
<b>Topic</b>	Particle & Nuclear Physics
<b>Sub Topic</b>	Nuclear Physics
<b>Paper Type</b>	Theory
<b>Booklet</b>	Question paper 3

**Time Allowed:** 81 minutes


**Score:** /67

**Percentage:** /100

CHEMISTRY ONLINE

A*	A	B	C	D	E	U
>85%	77.5%	70%	62.5%	57.5%	45%	<45%

- 1 One possible nuclear fission reaction is



Barium-141 ( ${}_{56}^{141}\text{Ba}$ ) and krypton-92 ( ${}_{36}^{92}\text{Kr}$ ) are both  $\beta$ -emitters.  
Barium-141 has a half-life of 18 minutes and a decay constant of  $6.4 \times 10^{-4} \text{ s}^{-1}$ .  
The half-life of krypton-92 is 3.0 seconds.

- (a) State what is meant by *decay constant*.

.....  
.....  
..... [2]

- (b) A mass of 1.2g of uranium-235 undergoes this nuclear reaction in a very short time (a few nanoseconds).

- (i) Calculate the number of barium-141 nuclei that are present immediately after the reaction has been completed.

number = ..... [2]


- (ii) Using your answer in (b)(i), calculate the total activity of the barium-141 and the krypton-92 a time of 1.0 hours after the fission reaction has taken place.

activity = ..... Bq [4]

- 2 (a) State what is meant by a *nuclear fusion reaction*.

.....  
.....  
.....[2]

- (b) One nuclear reaction that takes place in the core of the Sun is represented by the equation



Data for the nuclei are given in Fig. 8.1.

	mass/u
proton ${}^1_1\text{H}$	1.00728
deuterium ${}^2_1\text{H}$	2.01410
helium ${}^3_2\text{He}$	3.01605

**Fig. 8.1**

- (i) Calculate the energy, in joules, released in this reaction.

energy = ..... J [3]


- (ii) The temperature in the core of the Sun is approximately  $1.6 \times 10^7 \text{ K}$ .  
Suggest why such a high temperature is necessary for this reaction to take place.

.....  
.....  
.....[2]

- 3 (a) Explain why the mass of an  $\alpha$ -particle is less than the total mass of two individual protons and two individual neutrons.

.....  
 .....  
 ..... [2]

- (b) An equation for one possible nuclear reaction is



Data for the masses of the nuclei are given in Fig. 8.1.

		mass/u
proton	${}^1_1\text{p}$	1.00728
helium-4	${}^4_2\text{He}$	4.00260
nitrogen-14	${}^{14}_7\text{N}$	14.00307
oxygen-17	${}^{17}_8\text{O}$	16.99913

Fig. 8.1

- (i) Calculate the mass change, in u, associated with this reaction.

mass change = ..... u [2]

- (ii) Calculate the energy, in J, associated with the mass change in (i).

- (iii) Suggest and explain why, for this reaction to occur, the helium-4 nucleus must have a minimum speed.


.....

.....

..... [2]



- 4 When a neutron is captured by a uranium-235 nucleus, the outcome may be represented by the nuclear equation shown below.



- (a) (i) Use the equation to determine the value of x.

x = ..... [1]

- (ii) State the name of the particle represented by the symbol  ${}_{-1}^0\text{e}$ .

..... [1]

- (b) Some data for the nuclei in the reaction are given in Fig. 8.1.

		mass/u	binding energy per nucleon /MeV
uranium-235	( ${}_{92}^{235}\text{U}$ )	235.123	
molybdenum-95	( ${}_{42}^{95}\text{Mo}$ )	94.945	8.09
lanthanum-139	( ${}_{57}^{139}\text{La}$ )	138.955	7.92
proton	( ${}_1^1\text{p}$ )	1.007	
neutron	( ${}_0^1\text{n}$ )	1.009	

Fig. 8.1

Use data from Fig. 8.1 to

- (i) determine the binding energy, in u, of a nucleus of uranium-235,

binding energy = ..... u [3]

(ii) show that the binding energy per nucleon of a nucleus of uranium-235 is 7.18 MeV.

[3]

(c) The kinetic energy of the neutron before the reaction is negligible.  
Use data from (b) to calculate the total energy, in MeV, released in this reaction.

energy = ..... MeV [2]

CHEMISTRY ONLINE  
— TUITION —

- 5 (a) (i) State what is meant by the *decay constant* of a radioactive isotope.


.....  
.....  
..... [2]

- (ii) Show that the decay constant  $\lambda$  and the half-life  $t_{1/2}$  of an isotope are related by the expression

$$\lambda t_{1/2} = 0.693.$$

[3]

- (b) In order to determine the half-life of a sample of a radioactive isotope, a student measures the count rate near to the sample, as illustrated in Fig. 9.1.



**Fig. 9.1**



Initially, the measured count rate is 538 per minute. After a time of 8.0 hours, the measured count rate is 228 per minute.

Use these data to estimate the half-life of the isotope.

half-life = ..... hours [3]

- (c) The accepted value of the half-life of the isotope in (b) is 5.8 hours.  
The difference between this value for the half-life and that calculated in (b) cannot be explained by reference to faulty equipment.

Suggest two possible reasons for this difference.

1. ....  
.....  
2. ....  
.....

[2]

CHEMISTRY ONLINE  
— TUITION —

- 6 The element strontium has at least 16 isotopes. One of these isotopes is strontium-89. This isotope has a half-life of 52 days.

(a) State what is meant by *isotopes*.

.....  
.....  
..... [2]

(b) Calculate the probability per second of decay of a nucleus of strontium-89.

probability = ..... s<sup>-1</sup> [3]

(c) A laboratory prepares a strontium-89 source.

The activity of this source is measured 21 days after preparation of the source and is found to be  $7.4 \times 10^6$  Bq.

Determine, for the strontium-89 source at the time that it was prepared,

(i) the activity,

activity = ..... Bq [2]

(ii) the mass of strontium-89.

mass = ..... g [2]

- 7 The isotope phosphorus-33 ( $^{33}_{15}\text{P}$ ) undergoes  $\beta$ -decay to form sulfur-33 ( $^{33}_{16}\text{S}$ ), which is stable.  
The half-life of phosphorus-33 is 24.8 days.

(a) (i) Define radioactive *half-life*.

.....  
.....  
.....[2]

(ii) Show that the decay constant of phosphorus-33 is  $3.23 \times 10^{-7} \text{ s}^{-1}$ .

[1]

(b) A pure sample of phosphorus-33 has an initial activity of  $3.7 \times 10^6 \text{ Bq}$ .

Calculate

(i) the initial number of phosphorus-33 nuclei in the sample,

number = .....[2]

(ii) the number of phosphorus-33 nuclei remaining in the sample after 30 days.

number = .....[2]

- (c) After 30 days, the sample in (b) will contain phosphorus-33 and sulfur-33 nuclei.  
Use your answers in (b) to calculate the ratio

$$\frac{\text{number of phosphorus-33 nuclei after 30 days}}{\text{number of sulfur-33 nuclei after 30 days}} .$$

ratio = .....[2]

CHEMISTRY ONLINE  
— TUITION —

**8** Radon-222 is a radioactive element having a half-life of 3.82 days.

Radon-222, when found in atmospheric air, can present a health hazard. Safety measures should be taken when the activity of radon-222 exceeds 200 Bq per cubic metre of air.

**(a) (i)** Define radioactive *decay constant*.

.....  
.....  
..... [2]

**(ii)** Show that the decay constant of radon-222 is  $2.1 \times 10^{-6} \text{ s}^{-1}$ .

[1]

**(b)** A volume of  $1.0 \text{ m}^3$  of atmospheric air contains  $2.5 \times 10^{25}$  molecules.

Calculate the ratio

$$\frac{\text{number of air molecules in } 1.0 \text{ m}^3 \text{ of atmospheric air}}{\text{number of radon-222 atoms in } 1.0 \text{ m}^3 \text{ of atmospheric air}}$$

for the minimum activity of radon-222 at which safety measures should be taken.

ratio = ..... [3]